

**Child Protection Foundation  
Training**

# **Allegations of Sexual Abuse**

# Core Competencies


- ▶ Understand the definitions of child sexual abuse and the allegations
- ▶ Articulate the procedures for investigating allegations of child sexual abuse

# Definition


- ▶ “The involvement of children and adolescents in sexual activities they do not understand, to which they cannot give informed consent, or that violates social taboos.”

▮ Kempe, *Sexual Abuse: Another Hidden Pediatric Problem*, 1977


# Illinois Legal Definition

- ▶ **Sexual conduct - any intentional or knowing touching or fondling by the victim or the accused**, either directly or through clothing, of the sex organs, anus or breast of the victim or the accused, or any part of the body of a child under 13 years of age, for the purpose of sexual gratification or arousal
  - ▶ **Sexual penetration - any contact, however slight**, between the sex organ or anus of one person by an object, the sex organ, mouth or anus of another person, or **any intrusion, however slight**, of any part of the body of one person or object into the sex organ or anus of another person
- 

# Characteristics

- ▶ Forms of sexual abuse
    - Exhibitionism / voyeurism
    - Fondling / frottage
    - Vulvar coitus
    - Intercourse
    - Pornography
  - ▶ Psychological or physical coercion
  - ▶ Manipulation or exploitation
  - ▶ Misuse of position of authority / trust
- 

# Incidence


- ▶ Accounts for 10% of reported CA/N per year
  - ▶ Girls
    - About 20% of college-age women report some form of sexual abuse occurring before age 18
  - ▶ Boys
    - About 5-10% of college-age men report some form of sexual abuse occurring before age 18
  - ▶ Less than 10% of victims disclose at the time the abuse occurs
  - ▶ Children are most vulnerable between the ages of 7 and 13
- 

# Victims

- ▶ Victims: 80% female, 20% male
- ▶ Family “risk factors”
  - Single parent home, homes with step-parents
- ▶ Younger kids – more often victimized in home
- ▶ Older kids – more often victimized out of home
- ▶ It is estimated that each year there are about 400,000 children born who will become victims of child sexual abuse

# Perpetrators

- ▶ 90% are male
- ▶ 75 – 90% are known to victim
- ▶ 25 – 50% are relatives
- ▶ The younger the child victim, the more likely it is that the perpetrator is a juvenile. Juveniles are the offenders in 43% of assaults on children under age 6. Of these offenders, 14% are under the age of 12





**THERE ARE NO  
TYPICAL RISK  
FACTORS TO  
PROFILE  
PERPETRATORS**



# Presentation & Indicators

## ► Crisis

- Injury
- STD
- Pregnancy
- Drug overdose
- Suicide attempt
- Runaway

## ► Behavior problems

- Regression
- Sleep disorder
- School problems
- Depression
- Acting out
- Inappropriate sexual behavior



# Disclosure

- ▶ Most child victims never disclose
- ▶ Disclosure often delayed (18 months)
- ▶ Disclosure often incomplete
- ▶ Progression of disclosure:
  - Denial, disclosure, **recant**, reaffirmation

In Chicago alone, an estimated 3,200 incidents of child sex abuse are not reported each year. Abused kids are 59 percent more likely to be arrested, 66 percent more likely to use drugs, and 30 percent more likely to be abusive as adults.




**THE HISTORY IS  
OFTEN THE ONLY  
EVIDENCE OF  
SEXUAL ABUSE...**

# ...MOST PHYSICAL EXAMS ARE NORMAL

In the vast majority of cases where there is credible evidence that a child has been penetrated, only between 5 and 15% of those children will have genital injuries consistent with sexual abuse.

<https://victimsofcrime.org/media/reporting-on-child-sexual-abuse/child-sexual-abuse-statistics>

# Characteristics of Normal Sexual Behaviors

- ▶ Consenting
  - ▶ Developmentally appropriate
  - ▶ Mutually motivated by curiosity
  - ▶ Involves pleasurable feelings
  - ▶ Engage with peers of similar cognitive level
- 

# Normal Behaviors

- ▶ 2-3 y/o
  - Masturbation
  - Exhibitionism
  - Same/opposite sex exploration
- ▶ 6-7 y/o
  - Decreased overt sexual behaviors
  - Curious about sex, dirty words, etc.
- ▶ Puberty
  - Sexual activity with peers
  - 1/3 of 9<sup>th</sup> & 2/3 of 12<sup>th</sup> graders have had sex

# Characteristics of Abnormal Sexual Behaviors

- ▶ Coercion or pressure
- ▶ Developmentally inappropriate
- ▶ Motivated by one child's needs
  - Based on outside influences such as abuse
- ▶ Involve pain or discomfort
- ▶ Engages with non-peers
  - Older child on younger child




# Concerning Behaviors

- ▶ Touching adult genitals
- ▶ Making adults touch their genitals
- ▶ Masturbation with objects
- ▶ Masturbation causing pain
- ▶ Attempting or simulating intercourse
- ▶ Attempting or simulating oral sex

□ RM Reece & S Ludwig. *Child Abuse: Medical Diagnosis and Management*, Second edition, 2001



**REMEMBER...  
THESE BEHAVIORS  
ARE NOT DIAGNOSTIC  
AND SHOULD NEVER  
BE VIEWED IN  
ISOLATION...**




**...THESE BEHAVIORS  
MUST BE  
CORROBORATED  
WITH A HISTORY!**

# Medical Evaluation – History



## ▶ **HISTORY IS THE MOST IMPORTANT PART OF THE EVALUATION!**

- Interview child and caretaker separately
  - ▢ Interview should be in safe or neutral location
- Ask open-ended, non-leading questions
- Use patient's terminology for genital area
- Use direct quotes in the documentation

# Medical Evaluation - Physical Exam

- ▶ External visual genital exam in majority
  - ▶ No speculum used in prepubertal girls
  - ▶ **Complete** physical exam
  - ▶ Need for cultures determined case by case
  - ▶ Need for forensic kit determined case by case
- 

# Exam Positions





**What is the  
most common  
physical finding  
in sexual abuse  
evaluations?**

# **A Normal Exam!**

**75 - 80 % of  
exams are  
normal!**



# Why are most exams normal?

- ▶ Nature of sexual abuse
  - Fondling
  - Psychological rather than physical force
- ▶ Rapid healing of genital tissues
- ▶ Delay in seeking care
  - Disclosure often weeks or months after event
- ▶ Genital tissues are highly elastic
  - Tampons, masturbation, etc. do not cause traumatic injury to genital area

**What has  
been the  
response to  
sexual abuse?**

# Children's Advocacy Centers

- ▶ County facilities
- ▶ Comprehensive approach to services
  - **Offer collaboration of physicians, DCFS, police, state's attorneys**
  - Provide specialized forensic interviews
  - Advocate for the child and family
  - Provide education
  - Refer for necessary services




# Goal


**To ensure that children  
are not re-victimized by  
the very system  
designed to protect  
them**



# Coordinated Approach

- ▶ Interview of victim
  - ▶ Interview of all caretakers
  - ▶ Interview of alleged perpetrator
  - ▶ Medical evaluation
  - ▶ Medical records including
    - Documentation of history obtained
    - Documentation of physical exam
    - Culture results / lab tests
  - ▶ Police reports
- 


# Allegations

- ▶ #18 – Sexually Transmitted Diseases
  - ▶ #19 – Sexual Penetration
  - ▶ #20 – Sexual Exploitation
  - ▶ #21 – Sexual Molestation
  - ▶ #22 – Substantial Risk of Sexual Injury
- 

# #18-Sexually Transmitted Diseases

- ▶ “A disease which was acquired originally as a result of sexual penetration or sexual conduct with an individual who is afflicted with the disease.”

# STD List


- ▶ Chlamydia (*Chlamydia trachomatis*)
  - ▶ Gonorrhea (*Neisseria gonorrhea*; GC)
  - ▶ Genital warts (HPV; Human papilloma virus)
  - ▶ Syphilis (*Treponema pallidum*)
  - ▶ Genital herpes (HSV; Herpes simplex virus)
  - ▶ *Trichomonas vaginalis*
  - ▶ HIV / AIDS
  - ▶ Pubic lice
- 




# Background

- ▶ Statistics
  - 2-15% of sexually abused children
- ▶ Modes of transmission
  - At birth or in utero
  - Sexual contact
  - Casual contact?
- ▶ Clinical presentation
  - Discharge, rash, pain, bleeding
  - No symptoms

# Sexual Abuse Association

- ▶ **Gonorrhea**      ☐      **certain**
  - ▶ **Syphilis**      ☐      **certain**
  - ▶ **Chlamydia**      ☐      **probable**
  - ▶ **HPV**      ☐      **probable**
  - ▶ **Trichomonas**      ☐      **probable**
  - ▶ **HSV type 2**      ☐      **probable**
  - ▶ **HSV type 1**      ☐      **possible**
  - ▶ **HIV**      ☐      **uncertain**
- 

# **Gonorrhea, Chlamydia, & Syphilis**

- ▶ Not transmitted thru casual contact!
  - ▶ Not transmitted thru toilet seats!
  - ▶ Outside infancy, consider these transmitted through sexual contact
- 

# HPV (Warts)

- ▶ Difficult to determine etiology
- ▶ Incubation period unknown
  - Transmission from birth reported up to 2yo??
  - Large variation in when they are first detected
  - Typing of HPV not helpful
- ▶ In children > 2-3yo, HPV highly suspicious for sexual abuse


# HSV (Herpes)

- ▶ HSV Type 1 – oral (“cold sores”)
  - May transmit to self (autoinoculation)
  - Screen for sexual abuse (i.e. oral sex)
- ▶ HSV Type 2 – genital
  - Suspicious for sexual abuse
  - Unlikely but possible from non-sexual contact
- ▶ Diagnosis
  - Viral culture
  - Typing important but not diagnostic


# HIV

- ▶ Reported cases of HIV due to sexual abuse exist
- ▶ Incidence of sexual abuse transmission unknown
- ▶ Screen all sexually abused children for HIV
  - Positives may be from in utero transmission
  - Important to look at mother's prenatal screen
- ▶ Diagnosis from in utero transmission may be delayed as children may have no symptoms

# Procedure

- ▶ Forensic Interview
  - ▶ Re-exam by sexual abuse expert
  - ▶ Assess safety of environment
    - Removal if perpetrator in home
    - Safety plan if perpetrator unknown
  - ▶ Follow-up testing as recommended
  - ▶ Referral for sexual abuse counseling
  - ▶ Urgent medical exam for siblings / other children in the home
- 

# Medical Management

- ▶ Pregnancy test
  - ▶ Tests for other STD's
    - Genital swabs: GC, Chlamydia, HSV, Trichomonas
    - Blood tests: HIV, RPR (Syphilis), Hepatitis B
  - ▶ Antibiotics
  - ▶ **Forensic kit if contact < 72 hours**
  - ▶ **Pregnancy and HIV prophylaxis when needed**
- 




# STD testing of caretakers...

- ▶ If positive for STD's...
  - May link perpetrator to victim
  - Helpful evidence for prosecution
- ▶ If negative for STD's...
  - Does not eliminate perpetrator
    - ▢ May clear infection spontaneously
    - ▢ May be treated unknowingly
    - ▢ May conceal treatment intentionally


## #19 – Sexual Penetration

- ▶ “Any contact, however slight.....”
- ▶ “Any intrusion, however slight.....”

## #21 – Sexual Molestation

- “Sexual conduct [including fondling or inappropriate touching]... for arousal or gratification of sexual needs or desires.”
- 

# REMEMBER...

- ▶ The history is the most important evidence!
  - ▶ Physical exam is normal in 75-80%!
- 

# Procedures

- ▶ Forensic Interview
- ▶ Child needs emergent medical exam if
  - Last contact < 72 hours
  - Symptoms such as pain, bleeding, discharge
  - Forensic evidence collection kit needed
- ▶ Child needs non-emergent medical exam if
  - Last contact > 72 hours
  - No acute symptoms
  - Safe environment ensured


**REMEMBER...**

**THE CHILD NEEDS AN  
EXAM BY A  
PRACTITIONER WITH  
EXPERTISE IN CHILD  
SEXUAL ABUSE**

# The Role of the Medical Provider

- ▶ Gathering the medical history
- ▶ Evaluating the medical and mental health needs of the child
- ▶ Educating families, multidisciplinary team partners, judges, and jurors in the appropriate assessment, interpretation of findings and management of sexually abuse children and adolescents

Adams, J. et al. (2015) Updated Guidelines for the Medical Assessment and Care of Children who may have been Sexually Abused. (In Press) North American Society for Pediatric and Adolescent Gynecology.



# Location of Evaluation


- ▶ Use the ER only for emergency exams!
  - Chaotic, hurried environment
  - Not child-friendly atmosphere
  - Variable examiner knowledge / expertise
- ▶ CAC / sexual abuse expert for non-emergent exams!
  - Calm environment w/ adequate time allotment
  - Child friendly atmosphere
  - Examiner with expertise in sexual abuse

# Accidental Injury

- ▶ Straddle injury is most common
- ▶ Presents with clear history
- ▶ Primarily external injury
  - Bruises or cuts on the labia
  - Hymen rarely injured
- ▶ Presents with pain or bleeding



# #22 - Substantial Risk of Sexual Injury

- ▶ Sex offender has access to child
  - ▶ Siblings / other children in same household as alleged offender
  - ▶ Persistent, highly sexualized behavior or knowledge in a young child
- 

# Procedure

- ▶ Forensic interview if indicated
- ▶ Medical exam if forensic interview indicative of sexual abuse

# Case #1

- ▶ 5 year old female seen for vaginal discharge
- ▶ No disclosure from child
- ▶ No concerns from caretaker
- ▶ Physical exam normal
- ▶ Cultures positive for Gonorrhea
- ▶ Child called back to ED for treatment
- ▶ ED reports to DCFS and requests J-response

**WHAT ARE THE NEXT STEPS?**



# Case #2

- ▶ 8 year old male reported to DCFS hotline by teacher after disclosing that his uncle “touches [his] booty and [his] private area”
- ▶ Taken by police to the ED
- ▶ Child repeats disclosure to physician
- ▶ Exam is normal

**WHAT ARE THE NEXT STEPS?**



# Case #3

- ▶ 2 year old female reported to DCFS hotline by grandmother for suspicion that her “father is messing with her”
- ▶ GM’s suspicion based on crying when taken to dad’s home and sexually inappropriate behavior
- ▶ No interview / medical evaluation done yet

**WHAT ARE THE NEXT STEPS?**



# **Summary - Sexual Abuse Unit**

**Can you complete a sexual abuse investigation at this time?**

**What else do you need?**



# Questions?

